



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

BUFFALO, N. Y.

Passed Asst. Surg. Frost reports:

During the week ended October 12, 1912, 10 cases of poliomyelitis and 3 deaths therefrom were reported to the commissioner of health of Buffalo.

The total number of cases of poliomyelitis reported in Buffalo from January 1 to October 12, 1912, is 326. According to the best information at present available these cases may be classified as follows:

(1) Poliomyelitis with paralysis or paresis.....	276	
(2) Poliomyelitis, abortive	12	288
(3) Suspected poliomyelitis, diagnosis doubtful	11	
(4) Not poliomyelitis, erroneously reported.....	27	38
	<hr/> 326	<hr/> 326

The total number of deaths reported as due to poliomyelitis is 39. Of these, one was found at autopsy to have been due to tubercular meningitis; in three the cause of death is doubtful, leaving 35 deaths which may be considered with reasonable certainty as due to poliomyelitis.

WESTERN NEW YORK.

Reports have been received from the New York State Department of Health of 104 cases with 16 deaths in western New York, outside of Buffalo. These cases, with five exceptions, have occurred since July 1, 1912. They are distributed in Cattaraugus, Chautauqua, Erie, Genesee, Livingston, Niagara, Orleans, Wyoming, and Allegany Counties.

LOS ANGELES, CAL.

Surg. Brooks reports: A case of poliomyelitis, male, age 2 years, was reported in this city during the week ended October 5, 1912. No death from the disease occurred during the week. At the close of the week one case remained in the municipal hospital and four cases in quarantine in residences.

RABIES.

During the week ended September 28, 1912, a case of rabies was reported by the city health officer at Wilkinsburg, Pa.

TETANUS.

**CASES AND DEATHS REPORTED BY CITY HEALTH AUTHORITIES
FOR THE WEEK ENDED SEPT. 28, 1912.**

Place.	Cases.	Deaths.	Place.	Cases.	Deaths.
Bayonne, N. J.		1	Milwaukee, Wis.		1
Clinton, Mass.	1	1	New York, N. Y.		1
Los Angeles, Cal.	1		Pittsburgh, Pa.	1	